



Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Folicur® 3.6 F Foliar Fungicide

For Use on Asparagus to control Rust
(*Puccinia* spp.)

SECTION 18 EMERGENCY EXEMPTION
File Symbol: 04-MI-02

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MICHIGAN UNDER SECTION 18 EMERGENCY EXEMPTION

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Folicur® 3.6 F Foliar Fungicide (EPA Registration Number 264-752, formerly EPA Registration Number 3125-394).

Any adverse effects resulting from the use of Folicur® 3.6 F Foliar Fungicide under this emergency exemption must be immediately reported to the State of Michigan Department of Agriculture (517) 335-6542.

EFFECTIVE PERIOD:

This labeling becomes effective May 1, 2004 expires on November 1, 2004.

DIRECTIONS FOR USE

A maximum of 12,000 acres of asparagus in Michigan may be treated.

RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF FOLICUR® 3.6 F
Asparagus	Rust (<i>Puccinia</i> spp.)	4 – 6 fl. oz. per acre

Apply FOLICUR® 3.6 F as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl. oz. FOLICUR® 3.6 F per acre (0.11 lb ai – 0.17 lb ai per acre) when alternated with a protectant fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Apply with a spray adjuvant such as Induce.

RESTRICTIONS: Do not apply to harvestable spears. Do not apply within 180 days of harvest. Do not make more than 4 applications per growing season. Do not apply more than 16 fl. oz (0.45 lb ai) FOLICUR® 3.6 F per growing season.

GENERAL DIRECTIONS: Applications may be made using either ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. Apply the specified dosage in a minimum of 10 gallons of spray solution per acre by ground sprayer or a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)
or

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).